



US 20030099756A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0099756 A1**

Gow et al. (43) **Pub. Date: May 29, 2003**

(54) **METHOD OF PRODUCING A PROCESSED KAVA PRODUCT HAVING AN ALTERED KAVALACTONE DISTRIBUTION AND PROCESSED KAVA PRODUCTS PRODUCED USING THE SAME**

Publication Classification

(51) **Int. Cl.⁷** **A23K 1/00**
(52) **U.S. Cl.** **426/615**

(76) **Inventors: Robert Gow, Naples, FL (US); John Pierce, Moreno Valley, CA (US); Brian Pierce, Thousand Oaks, CA (US); William Birdsall, Naples, FL (US)**

(57) **ABSTRACT**

A method of producing a processed kava product involves using an extraction solvent, such as liquid CO₂, to preferentially extract different kavalactones from the source material at different rates. By controlling the extraction parameters and stopping the extraction before all of the kavalactones have been extracted or allowing the extracted kavalactones to be preferentially precipitated in one or more collection environments, a processed kava product can be produced that has a kavalactone distribution profile that can differ substantially from that of the source material. As a result, roots from a less desirable kava cultivar can be used to produce a processed kava product which has a kavalactone distribution profile that is similar to that of a highly desired cultivar. The kava paste can be further processed to produce a dry flowable powder suitable for use in, e.g., a tableting formula. A rapid dissolve tablet formulation for use in the delivery of kavalactones is also disclosed.

Correspondence Address:
CLIFFORD CHANCE US LLP
200 PARK AVENUE
NEW YORK, NY 10166 (US)

(21) **Appl. No.: 10/263,579**

(22) **Filed: Oct. 3, 2002**

Related U.S. Application Data

(60) **Provisional application No. 60/326,928, filed on Oct. 3, 2001. Provisional application No. 60/369,889, filed on Apr. 3, 2002.**




Fig. 1





Fig. 2

Average DHK/K and Y/DMY Kavalactone Concentration Ratios Measured in the Lateral Roots for Five Usage Classes of Vanuatu Kava Cultivars: Daily, Custom, Two-day, Medicinal and No Drink; Fiji Lateral Roots; and Two Kava Extracts: Alden Botanica Extract and Laitan Extract



Yangonin / Desmethoxyyangonin (dB)

Fig. 3




FIG. 4




FIG. 5




FIG. 6

Liquid CO2 Extract of Lateral Roots of Hawaiian Kava Cultivar: (DHK%/K%) vs. (Y%/DMY%) in dB Units




FIG. 7

METHOD OF PRODUCING A PROCESSED KAVA PRODUCT HAVING AN ALTERED KAVALACTONE DISTRIBUTION AND PROCESSED KAVA PRODUCTS PRODUCED USING THE SAME

CROSS-REFERENCE TO RELATED APPLICATIONS:

[0001] This application is claims priority under 35 U.S.C. § 119 to U.S. Provisional Patent Application Serial No. 60/326,928 entitled "Methods For Processing Kava Root (Piper Methysticum) And Kava-Derived Products Generated By These Methods Having Controllable Kavalactone Ratios", filed on Oct. 3, 2001, and U.S. Provisional Application Serial No. 60/369,889 entitled "Dry Flowable Kava Extract Powder And Fast-Dissolve Kava Extract Composition For Oral Delivery", filed on Apr. 3, 2002. The entire contents of both of these applications are hereby expressly incorporated by reference.

FIELD OF THE INVENTION(S)

[0002] This invention is related to methods for processing kava to produce a kava extract with particular kavalactone concentration profile, a processed kava product, such as a dry flowable kava extract powder, and the use of such a powder in a rapid-dissolve tablet for oral delivery.

BACKGROUND

[0003] Kava plants, a type of pepper plant also known as Piper methysticum, are generally found in Polynesia, Melanesia, and Micronesia. The kava plant contains high concentrations of kavalactones (sometimes referred to as kavapyrones), including kavain, methysticin, yangonin, dihydromethysticin, desmethoxyyangonin, and dihydrokavain, and has been used as an herbal medicine. The prized part of the kava plant is the root system because it contains the highest concentrations of the active kavalactones. Kavalactones are also found in other parts of the plant. General information about kava plants can be found in Vincent Lebot, et al., "Kava The Pacific Elixir, The Definitive Guide to its Ethnobotany, History, and Chemistry", Inner Traditions Intl. Ltd. (March 1997).

[0004] After a kava plant is harvested, the root is conventionally processed to generate a consumable product. In conventional and historical methods, the root is dried and ground to a powder. This powder contains not only the kavalactones but also plant oils, resins, and other substances. The powder can be mixed with water to form a beverage. The resinous ground dried root can also be packaged into capsules or other forms of delivery.

[0005] The effects of ingesting kava root extract will vary from person to person. Common effects include a state of relaxation and a reduction in muscle tenseness. Kava root has also been used to help sufferers of insomnia. It can also produce a mild state of euphoria. The ratios of the various kavalactones in the consumed product has a large impact on the effect experienced by the user.

[0006] There are a large number of different kava cultivars, each of which has differing ratios of the different kavalactones. Through historical knowledge and experimentation, various kava cultivars have been classified according to their effect, such as "daily" or "one-day", "custom",

"two-day", "medicinal", and "no drink". "Daily" kava cultivars are generally consumed daily and have relatively short-term effects. "Custom" kava cultivars tend to be used for ritual and ceremonial purposes. "Two-day" kava cultivars tend to have pronounced physiological effects and are known as "two day" because the drinker is usually affected for two days. "Medicinal" kava cultivars are specific for the treatment of ailments like rheumatism. "No Drink" kava cultivars are not recommended for consumption because they tend to induce nausea and other undesirable effects.

[0007] More recently, there have been concerns associating kava with occurrences of liver damage, although these reports are primarily anecdotal. Studies done in the 1970s and 1980s have found that "two day" kava cultivars have particularly high concentrations of methysticin and dihydromethysticin. These kavalactones contain a methylenedioxyphenyl functional group which, when activated through metabolic activity, forms intermediary compounds that can inactivate multiple P450 enzymes. There are some suggestions that inhibition of P450 enzymes can interfere with the metabolism of many pharmaceuticals. The solution proposed in the prior art is the development of new kava cultivars with reduced amounts of methysticin and dihydromethysticin and increased kavain content.

[0008] One of the most highly prized kava cultivars is a one-day cultivar that grows in Vanuatu. Relative to other cultivars, this cultivar is naturally low in methysticin and dihydromethysticin and has a proportionally higher concentration of kavain. However, this cultivar is not widely grown or available and most commercially available cultivars have increased levels of methysticin and dihydromethysticin and lower concentrations of kavain. Although breeding of new cultivars that can be widely grown and which have lower concentrations of methysticin and dihydromethysticin and increased concentrations of kavain is possible, this can be a time consuming process and is not assured of success. In addition, the kavalactones suspected of producing toxic effects when consumed by humans are also toxic to various insects and microorganisms, such as fungi. Thus, the hardest cultivars are generally those which are least suitable for human consumption.

[0009] In a kava extraction process, the rate at which various kavalactone components are extracted varies according to the extraction solvent and extraction conditions used. However, kava root extracts are conventionally sold based upon total kavalactone content. Thus, the focus of conventional commercial extraction techniques has been to extract as much of the kavalactone content from the root source as possible and knowledge regarding different rates of extraction for different kavalactones is used to adjust the extraction process or select appropriate extraction steps to provide for the greatest total kavalactone extraction.

[0010] Various extraction techniques have been used to process kava root. Such techniques include the use of supercritical carbon dioxide (CO₂) extraction, fluorocarbon extraction, and the use of various other organic and non-organic solvents to remove the kavalactones from the dried root. The resulting bulk extract is in the form of a paste.

[0011] Supercritical CO₂ processes have been found to extract the largest quantity of kavalactones from the root and thus conventional practice has been to focus on the use of supercritical CO₂ to extract all of the kavalactones from the

root in the production of kava extracts. As will be appreciated, bulk extracts of substantially all of the kavalactones in the root will generally contain kavalactones in the same or nearly the same proportions as present in the source root. Because of the differing effects produced by variations in the kavalactone ratios of different cultivars, producers of such kava products are limited to specific kava cultivars as source materials in order to produce a kava product that produces a desired effect.

[0012] To the extent that the distribution profile of the various kavalactones in the extract have been of concern, conventional practice is to use additional processing steps, such as high pressure liquid chromatography (HPLC), to extract individual kavalactones. For example, HPLC and other chromatographic techniques, such as "flash" chromatography, have been used to process bulk kava extracts to isolate individual kavalactones. This process is cumbersome to implement and requires the use of machinery that is bulky and expensive to operate. Methods for extracting and purifying kava, including supercritical CO₂ extraction and chromatography, are discussed in PCT publication WO 00/72861 to Martin et al. entitled "Pharmaceutical Preparations of Bioactive Substances Extracted from Natural Sources." It would be advantageous and is an object of the present invention to provide a method of processing kava to provide an extract of kava as a paste, powder, or in other forms, and which allows the kavalactone profile to be altered during processing so as to have reduced levels of methysticin and dihydromethysticin and increased levels of kavain and dihydrokavain. It would be additionally advantageous if such techniques allowed the production of a processed kava product that had a kavalactone profile similar to (or improved upon) that of the most preferred "one-day" cultivars from Vanuatu by using a kava feedstock from a less desired cultivar. There would be yet a further advantage if such an altered-profile kava product could be produced using a minimal number of processing steps and without requiring additional processing, such as via chromatography, to isolate each individual kavalactone from the extract and then recombine the isolated kavalactones as appropriate.

[0013] There has been widespread use of bulk kava paste extracts as dietary supplements, e.g., mixed in drinks or packaged in capsules. Because kava paste is not always well suited for processing into consumable form or for distribution, e.g., as a dietary supplement, it would also be advantageous to produce a dry flowable kava powder. The powder can be distributed as is or further processed, e.g., to produce kava tablets.

[0014] When consumed, kava is typically swallowed. The kavalactones then pass through the stomach and intestines as they are absorbed into the bloodstream. It can take a relatively long time for the body to process kava and the effect of the ingested kavalactones can then be further diminished as the compounds are processed by the liver. It would therefore also be advantageous to deliver kava in the form of a rapid-dissolve tablet which would permit the kavalactones to be absorbed orally.

SUMMARY OF THE INVENTION(S)

[0015] These and other objectives are met by use of the present methods of processing kava to produce a kava product, such as a paste, resin, oil, or a powder suitable for

use in a fast dissolve tablet and other applications, and which as a kavalactone profile that is altered from the profile of the kava feedstock. The present methodologies also allow the distribution of kavalactones to be altered or adjusted to achieve a particular kavalactone profile wherein the effectiveness of the kava can be enhanced or the negative effects reduced. Various other novel kava extract compositions are also presented.

[0016] Preferably, a two stage process is used. First, the kavalactones are extracted from the kava root material to form a kava paste. Preferably, the paste is generated using liquid CO₂ extraction techniques. However, under some conditions, supercritical CO₂ extraction techniques can be used. During processing, an initial pressurization/depressurization step can also be included to increase the rate at which the kavalactones can be extracted from the rootstock. The output of the first processing step is a viscous paste, oil, and/or resin.

[0017] The paste can then be subjected to a second or further refinement stages, if desired, to further purify the kavalactones and produce a highly refined paste extract. Preferably, the parameters of the second processing stage (and possibly the first stage as well) are selected to alter the kavalactone distribution in the raw material in order to achieve a predetermined distribution and concentration of the various kavalactones without the need of chromatography or other techniques conventionally used to extract individual kavalactones.

[0018] Advantageously, the use of the extraction techniques according to the various embodiments of the invention thus allows high quality kavalactone having a desired profile, such as reduced levels of methysticin and dihydromethysticin and increased levels of kavain and dihydrokavain, to be easily and economically produced from a kava feedstock, such as dried ground root, which has an undesired kavalactone profile. In a particular example, an undesired kava stock, such as a "two-day" or "no-drink" cultivar, can be processed to produce a kava extract having a kavalactone distribution profile which is comparable to the most desired "one-day" Vanuatu cultivar.

[0019] For example, processing of kava using CO₂ at pressures of 1100 psi or greater and at a temperature greater than 31° C. places the CO₂ in a supercritical state and will extract on the order of 80% to 100% of the kavalactones in the source. However, the extraction rate of the various kavalactones are fairly similar and the resulting extract will have a kavalactone profile that is similar to, if not identical to, that of the feedstock. (However, with careful control over the extraction parameters, subtle changes in the kavalactone profile can be made using supercritical CO₂ extraction.)

[0020] In contrast, it has been found that processing the kava feedstock using liquid CO₂ at pressures of between about 1100 psi to about 1800 psi and temperatures between about 0° C. and about 25° C. produces preferential extraction wherein methysticin and dihydromethysticin are extracted at a considerably slower rate than other kavalactones. It has been further found that processing at between about 1100 to about 1500 psi at temperatures between about 5° C. and about 10° C. also produces an increased relative concentrations of kavain and dihydrokavain relative to other kavalactones in the extract.

[0021] Through use of the present techniques, kava extracts can be produced having a kavalactone profile that is

unlike those found in nature and the profile can be tailored to meet particular design considerations. Preferably, the kava product has kavalactone properties that have been selected to be low in the combined weight percentage of methysticin (M) and dihydromethysticin (DHM), low in the combined weight percentage of desmethoxy yangonin (DMY) and yangonin (Y), and high in the combined weight percentage of kavain (K) and dihydrokavain (DHK). The preferred kava product can be characterized according to one or more of the following attributes: (a) a combined weight percentage of the six major kavalactones methysticin (M), dihydromethysticin (DHM), yangonin (Y), desmethoxyyangonin (DMY), kavain (K), and dihydrokavain (DHK) of between about 20% to a maximum of about 90%; (b) a combined weight percentage of M and DHM between about 5-15% to about 29%; (c) a ratio of Y to DMY by weight, expressed as the logarithmic function $10 \cdot \text{LOG}_{10}(Y/DMY)$ in dB units, from about -1 to about 2; (d) a ratio of DHK to K by weight, expressed as the logarithmic function $10 \cdot \text{LOG}_{10}(DHK/K)$ in dB units, from about -4 to about 1; (e) a combined weight percentage of Y and DMY between about 5% to about 20-25%, and (f) a combined weight percentage of flavokavain A and flavokavain B of between about 0.3% to about 3%.

[0022] Once a suitable kava paste, resin, or oil, is provided, it can then be processed for various uses. For direct ingestion, for example, a paste can be sweetened and flavored. In addition or alternatively, the paste can be mixed with other dietary supplements, such as the very sweet tasting herb Stevia, or other herbal or non-herbal ingredients.

[0023] A kava paste can also be further processed to prepare a high quality dry flowable powder which can be used, for example, to produce an ingestible kava tablet. In a first embodiment, a kava paste is combined with a carrier, such as maltodextrin, dextrose, or starches and mixed with a suitable solvent, such as ethyl alcohol or water. The mixture is then spray dried to produce a powder having grains comprised of kava extract and the carrier. In a second embodiment, an emulsion of kava paste is formed in water or ethyl alcohol using, e.g., magnesium carbonate, magnesium carbonate+silica (at up to about 2% by weight), whey protein, maltodextrin, carboxymethylcellulose and/or other suitable materials. The emulsion is then dried and powdered. In a third embodiment, the extract is processed with supercritical CO₂ and a carrier, such as maltodextrin, dextrose, or starches and then subjected to a rapid decompression of the supercritical fluid. As the fluid evaporates, the extract and kava are deposited as micro-sized particles.

[0024] The resulting powder, preferably having a preselected ratio of kavalactones, can then be formed into a tablet that, when placed in the mouth, dissolves rapidly over a period of between about 5 to about 180 seconds and preferably in about 15 to about 60 seconds. A tableting powder can be formed by combining between about 25% to about 60% by weight of the powdered kava extract with between about 30% to about 60% by weight of a dry water-soluble absorbant such as magnesium carbonate, or a diluent, such as lactose. Other dry tablet additives, such as one or more of a sweetener, flavoring and/or coloring agents, a binder, such as acacia or gum arabic, a lubricant, a disintegrant, and a

buffer, can also be added to the tableting powder. Preferably, the dry ingredients are screened to a particle size of between about 80 to about 100 mesh.

[0025] To form the tablets, the tableting powder is then wet with a solvent, such as alcohol or water, until it achieves a doughy consistency. The paste is then molded or pressed into tablets which are dried and packaged. The tablets can be of any suitable size, shape, and weight. Preferably, the tablets are formed into disks or wafers having a diameter of between about $\frac{1}{8}$ and about $\frac{1}{2}$ inch, a thickness of between about 0.08 and about 0.2 inch, and a weight of between about 50 mg to about 1000 mg. The tablets are preferably homogeneous. However, they can also be formed of regions of kava extract composition separated by non-kava extract regions in periodic or non-periodic sequences as may be desired.

BRIEF DESCRIPTION OF THE DRAWINGS

[0026] The following detailed description of the preferred embodiment(s) will be better understood with reference to the following figures in which:

[0027] FIG. 1 is a plot of total kavalactone weight percentages measured in the lateral roots for five usage classes of Vanuatu kava cultivars: Daily, Custom, Two-day, Medicinal and No Drink;

[0028] FIG. 2 is a plot of Methysticin+dihydromethysticin weight percentages measured in the lateral roots for five usage classes of Vanuatu kava cultivars;

[0029] FIG. 3 is a plot of average dihydrokavain+kavain vs. yangonin+desmethoxyyangonin kavalactone concentration ratios measured in the lateral roots for various kava cultivars and extracts; and

[0030] FIGS. 4-7 show plots of relative kavalactone concentrations for different kava cultivars and show changes in the kavalactone profile introduced during processing according to the present invention.

DESCRIPTION OF THE PREFERRED EMBODIMENT(S)

[0031] To better appreciate the utility of the present invention, it is useful to present data describing the differences between the total and relative amounts of various kavalactones in a variety of kava cultivars. The HPLC measurements for kava cultivars are taken from V. Lebot's and J. Levesque's paper in *Allertonia* (1989) entitled, "On The Origin And Distribution Of Kava (*piper methysticum* forst. F., *piperaceae*): A Phytochemical Approach". Lebot's and Levesque's procedure for the extraction and analysis of total kavalactone weight percentages and kavalactone concentration spectra for kava cultivars consisted of the following steps: First, select only the lateral roots of a kava plant that reached the age of two years. Next, dry lateral roots in an oven at 60-80° C. for eight hours, and then grind the roots into powder. Then, perform a six hour Soxhlet extraction of powdered root using chloroform as the solvent. Finally, analyze the chloroform extract using HPLC.

[0032] Note that Lebot and Levesque used only the lateral roots to measure the kavalactone concentration spectrum for a given kava cultivar. This is important because different parts of the kava plant (e.g., rhizome, stem, leaves) have

different kavalactone concentration spectra, as well as different total kavalactone weight percentages. Total kavalactone weight percentage is highest in the lateral roots, and decreases towards the aerial parts of the plant (pg. 67, Lebot, Merlin, Lindstrom, *Kava-The Pacific Elixir*, 1997). Lebot (1988) found in Vanuatu kava cultivars that when the total kavalactone weight percentage was about 15% in the lateral roots, it dropped to about 10% and 5% in the rhizome and basal stem, respectively. Lebot, et al (1997) on pp. 74-75 report that K and DMY are the dominant kavalactones in the rhizome, whereas DHK and DHM tend to be the kavalactones with the highest concentrations in the leaves. In summary, the kavalactone concentration spectrum and total kavalactone weight percentage for a given kava cultivar can be determined by a number of factors, which include age of the plant, method of extraction, as well as the part of the plant.

[0033] 66 of the 67 Vanuatu cultivars identified in Table 2 in Lebot and Levesque (1989) were analyzed (33 Daily, 11 Custom, 14 Two-day, 4 Medicinal, and 4 No Drink cultivars). Also analyzed were 26 of the 27 Fijian kava cultivars from Table 6 in Lebot and Levesque (1989). Lebot and Levesque did not classify these cultivars according to the system described above. HPLC was used to determine the total kavalactone weight percentages and kavalactone concentration spectra for the various samples.

[0034] FIG. 1 presents a plot of the total kavalactone weight percentages measured by Lebot and Levesque (1989) in the lateral roots for the five usage classes of Vanuatu kava cultivars: Daily (D), Custom (C), Two-day (T), Medicinal (M), and No Drink (N). As will be appreciated, the total kavalactone weight percentage does not uniquely identify the type of kava cultivar, e.g., there are several D, C, T and M kava cultivars with total kavalactone weight percentages in the range of 11-12%. FIG. 1 also shows that the selection of a kava cultivar with a very low total kavalactone percentage does not mean that it has the least effect because the class of No Drink kava cultivars has the lowest average total kavalactone concentration. Accordingly, kava cultivars that have high concentrations of desirable kavalactones may also contain high concentrations of undesirable kavalactones, making these cultivars unsuitable for use in producing consumable kava products without first altering the profile of the kavalactones.

[0035] FIG. 2 is a bar graph plotting number of Vanuatu kava cultivars vs. the combined M+DHM weight percentage measured by Lebot and Levesque (1989) in the lateral roots for the D, C, T, M and N classes of kava cultivars described for FIG. 1. Table 1 below summarizes the M%+DHM% averages for the five usage classes:

TABLE 1

Average Weight Percentage of Methysticin and Dihydromethysticin in Lateral Roots for the Five Usage Classes of Vanuatu Kava cultivars	
Kava Cultivar Usage Class	Average M% + DHM%
Daily	25 ± 4%
Custom	33 ± 9%
Two-day	39 ± 6%

TABLE 1-continued

Average Weight Percentage of Methysticin and Dihydromethysticin in Lateral Roots for the Five Usage Classes of Vanuatu Kava cultivars	
Kava Cultivar Usage Class	Average M% + DHM%
Medicinal	33 ± 13%
No Drink	57 ± 10%

[0036] FIG. 2 and Table 1 indicate that the frequency of use for a kava cultivar is proportional to the M%+DHM%, with the Daily kavas having the lowest average of 25±4%, the Two-day kavas are 14% higher at 39±6%, and the No Drink kavas have the highest M%+DHM% at 57±10%. Thus, the higher the M%+DHM% for a Vanuatu kava cultivar, the less frequent its use.

[0037] Moreover, M%+DHM% has been found to be proportional to the degree of induced nausea, which could help explain why frequency of use drops with increasing M%+DHM%. In particular, Lebot, et al (1997) report on pg. 78 that DHM is a major component of Two-day (or tudei) kavas that frequently produce nausea. According to the work of the Meyer group in Freiburg, the M and DHM kavalactones are absorbed the slowest in the gastrointestinal tract. The strong anticonvulsive and muscle relaxant action of M and DHM (see Meyer group work) combined with the fact that M and DHM remain for longer periods of time in the gastrointestinal tract are consistent with the induction of nausea resulting from kava preparations containing high percentages of M and DHM.

[0038] The relationships between M%+DHM% and Vanuatu kava cultivar usage presented in FIG. 2 and table 1 above are important steps in classifying Vanuatu kava cultivars, and can also be used to classify other kava preparations and products. However, this classification does not help with distinguishing the neuroactive effects of the different kava cultivars. In order to better distinguish the neuroactive effects of kava products, a two-dimensional logarithmic coordinate system is presented which maps the concentrations of the neuroactive kavalactones that cross the blood-brain barrier the fastest: K, DHK, Y and DMY. The Y/DMY ratio defines the "X-axis" for this coordinate system and the DHK/K ratio defines the "Y-axis". The logarithmic nature of the coordinate system comes from the fact that the DHK/K and Y/DMY concentration ratios are in dB's, which means $10 \cdot \text{LOG}_{10}(\text{DHK}/\text{K})$ and $10 \cdot \text{LOG}_{10}(\text{Y}/\text{DMY})$. A point centered on the origin (0,0) signifies that the DHK/K and Y/DMY ratios are both 1. Positive values reflect ratios greater than 1, and negative values correspond to ratios less than 1.

[0039] FIG. 3 is a map of average DHK/K vs. Y/DMY concentration ratios (in dB) for 1) lateral roots of the five different usage classes of Vanuatu kava cultivars, 2) lateral roots of 26 Fijian kava cultivars, 3) Alden Botanica's Essence of Kava (or EoK) extract of Vanuatu kava, and 4) Schwabe's Laitan kava extract. The DHK/K vs. Y/DMY map in FIG. 3 is effective in separating the five classes of Vanuatu kava cultivars, the Fijian kava cultivars, EoK, and Laitan. The average for the Fijian kavas falls in the fourth quadrant (DHK<1 and Y/DMY>1), in contrast to the averages for the D, C, M, and T Vanuatu kava cultivars in the first

quadrant (DHK>1 and Y/DMY>1), and the average for the N Vanuatu kava cultivars in the second quadrant (DHK>1 and Y/DMY<1). Although EoK is made from Vanuatu kava, the composition is about 20% lateral roots and 80% rhizome. The high percentage of rhizome in EoK could explain why its DHK/K ratio is less than 1 because K tends to dominate in the rhizome. The M%+DHM% for EoK is 21%, which is characteristic of a Vanuatu Daily kava. The lateral root and rhizome compositions in Laitan are not known, and so it is difficult to suggest a reason for its DHK/K vs. Y/DMY mapping. However, its M%+DHM% of 34% is closer to the average M%+DHM% for Vanuatu Custom, Medicinal and/ or Two-day kava cultivars and the average for Fijian kava cultivars. The high M%+DHM% of Laitan relative to that for EoK implies that Laitan will have a more muscle relaxing effect and a higher probability of inducing nausea.

[0040] As will be appreciated, there are many kava cultivars that contain kavalactone distributions which are not well suited for consumption. In order to generate a kava product, such as a paste or a dry flowable powder for use in fast dissolve tablet and other applications, raw kava root is processed to produce a kava paste, oil or resin (collectively referred to herein as "paste") that contains the kavalactones in the desired proportions, such as a distribution profile that corresponds to the most preferred Vanuatu one day cultivars. The paste can then be further processed, e.g., to produce a refined paste and/or a flowable powder. Preferred techniques are discussed below. These products generally can be referred to as a processed kava products.

[0041] As an initial step, the kava itself must be harvested, dried, and ground to size. The active kavalactones are most prevalent in the lateral roots of the plant. Various different cultivars are conventionally used, including but not limited to Isa, Mahakea, Moi, Papa Eleele, Borogu, Ahine, Kelai, Palisi, Puna Green, Hanakapiai, Hiwa, Spotted Hiwa, Nene, SIG, Apu Awa, Alia, Melomelo, and Pia. Each cultivar has a typical concentration of the various kavalactones resulting in different effects from ingesting the root or a root extract. The harvested roots are preferably dried to a moisture content of between 3% and 10%, most preferably around 5%, and then ground to a particle size suitable for subsequent processing. A typical particle size is between about 60 mesh to about 300 mesh, which correlates to about 250 μ to about 50 μ , respectively, and preferably to between about 100 μ and about 75 μ .

[0042] The powdered kava root is then processed to produce a kava paste using techniques described below. In a preferred technique, the kava is processed using CO₂ in the liquid phase, but supercritical phase CO₂ extraction can be used. A solvent modifier, such as ethyl alcohol or refrigerant chemicals, can be added. Preferably, prior to the primary extraction steps, the feedstock is initially processed by subjecting it to a high pressure carbon dioxide environment followed by a rapid decompression. This compression-decompression process advantageously results in a substantially higher kavalactone yield than achieved without the decompression step.

[0043] The paste can then be further processed to produce a flowable powder. Although a paste made according to the present invention is preferably used, kava pastes produced using various alternative techniques can alternatively be used. Such methods include those disclosed in Published

PCT Application WO 00/72861 to Martin et al. entitled "Pharmaceutical Preparations of Bioactive Substances Extracted from Natural Sources" and in U.S. Pat. No. 5,512,285 to Wilde entitled "Fragrance Extraction," the entire contents of which are both expressly incorporated by reference.

[0044] In a specific implementation of the method for producing the kava paste, the powdered dried root is placed in a stainless steel vessel and preprocessed by pressurizing the vessel with CO₂ to between about 1100 psi and about 4500 psi, preferably at least about 1800 psi, at a temperature of between about 10° C. and about 60° C., and preferably between about 15° C. and about 20° C. It is not required that the pressures reach the critical point of CO₂ (1100 psi and 31° C.) during this stage. After the desired pressure is reached and held for a period of time, ranging from continuous flow to fill and hold, where the hold time can vary from between about 5 minutes to about 60 minutes, the pressure in the extraction vessel is rapidly and substantially reduced, preferably to a level at or near one atmosphere. This preprocessing, while not necessary, allows for a higher efficiency of extraction in the subsequent processing steps.

[0045] As noted above, various extraction steps can be used to prepare the refined kava paste from a starting material. The particular techniques and solvents used are selected, to some extent, based on the moisture content present in the feedstock and on user preference. A specific factor that should be considered is the relative concentrations of the various kavalactones that is desired in the final product. For example, it can be beneficial to substantially reduce the amount of particular kavalactones in the extract which may cause interaction with various prescription drugs or cause various negative effects, while retaining high relative percentages of other kavalactones.

[0046] After the preferred decompression processing step is performed, the root contained within the stainless steel vessel is then processed to produce the kava extract paste. Various extraction processes (with or without the decompressive preprocessing) can preferably be used, with liquid CO₂ being the most preferred technique. The processes are preferably used in the alternative but two or more processes could be combined if desired. The processes are:

[0047] (1) CO₂ in the supercritical fluid phase.

[0048] (2) CO₂ in the liquid phase.

[0049] (3) CO₂ in the supercritical phase in combination with added ethyl alcohol as a modifier which is added in the range of 2% to 15%, and preferably between 2% and 10% based on the total mass.

[0050] (4) CO₂ in the liquid phase in combination with ethyl alcohol as a modifier which is added in the range of 2% to 15% and preferably between 2% and 10% based on the total mass.

[0051] (5) Refrigerant chemicals such as hydrofluorocarbons (HFCs), hydrochlorofluorocarbons (HCFCs), and/or chlorofluorocarbons (CFCs) such as: HFC-23, HFC-32, HFC-125, HFC-134a, HFC-143a, HFC-152a, R-404a, R-407c, R-410a, HCFC-22, HCFC-123, HCFC-141b, HCFC-142b, R-502, R-11, R-12, and R-113. A more complete listing of suitable refrigerant chemicals is in Table 2. These refrigerants are widely available from various chemical manufacturers and suppliers.

[0052] The selection of one extraction procedure over another depends on the objectives of the extraction, the kavalactone concentration profile of the raw starting material and the desired concentration profile of the extract end product. In order to extract the largest quantity of kavalactones in the aggregate, supercritical CO₂ extraction generally provides the best results. To alter or adjust the kavalactone profile, liquid CO₂ extraction techniques are more suitable. However, a carefully controlled supercritical CO₂ process may also be used. A gaseous CO₂ process may also be suitable under certain circumstances.

[0053] For example, if a starting kava feedstock has a kavalactone concentration profile (specified as weight percentages of the individual kavalactones) that is relatively low in (K+DHK) and relatively high in both (M+DHM) and (DMY+Y), and a kava product is desired that is relatively high in (K+DHK) and relatively low in both (M+DHM) and (DMY+Y), then the CO₂ extraction in the liquid phase becomes the preferred process. The figures discussed below illustrate how liquid CO₂ extraction transformed the kavalactone concentration profile of the starting materials. In summary, the above extraction procedures provide maximum flexibility in starting with any kava raw product to obtain and extract end product with a specific kavalactone concentration profile.

[0054] In a first embodiment of the process, CO₂ is used in the supercritical phase to perform the extraction step. The pressure is held at a pressure and temperature regime that between about 1100 psi and about 8000 psi and at a temperature between about 31° C. and about 80° C. During the process the extractable material is collected within a stainless steel collection vessel that could be the same one as that used in the first "low pressure" extraction step, and the supercritical CO₂ can either be recycled for future use or vented into the atmosphere. The solvent-to-feed ratio increases as the pressure and temperature of the supercritical CO₂ is lowered. It has been found that when the material is processed at a pressure of about 8000 psi and a temperature of about 80° C., all extractable material is obtained after a solvent-to-feed ratio of 8 is achieved.

[0055] In a second, most preferred, embodiment, CO₂ in the liquid phase is used to perform the extraction step. Since the CO₂ is in the liquid phase, the extraction temperature can not exceed about 31° C., however the pressure can rise to about 8000 psi. Conversely, the pressure can be maintained below about 1100 psi and the temperature can be varied from about 5° C. up to about 70° C. During this process the extractable material is collected within the stainless steel collection vessel (which could be the same one as that used in a first "low pressure" extraction step). The liquid CO₂ can either be recycled for future use or vented into the atmosphere.

[0056] In a particular example of liquid CO₂ extraction and fractionation, CO₂ at a pressure between about 1100 and about 1800 psi is used at a temperature of between about 5° C. and about 20° C. The low temperature during extraction is controlled via a recirculating chiller plumbed to jacketed extractors, and the pressure within the extraction vessel is controlled via a back pressure regulator. Because of preferential absorption, and as discussed more fully below with respect to FIGS. 4-7, the kavalactone profile of the dissolved kava can be substantially different from the profile of the source kava.

[0057] The extract-laden fluid emanating from the extractor is then collected. A single collection vessel can be used or the fluid can be sequentially processed using multiple collecting vessels to allow for further refinement of the kava extract. In a particular example, the liquid is first heated, e.g., by using a heat exchanger, to a temperature between about 30° C. and about 37° C. The vessel into which the fluid is deposited is also maintained at the same temperature. A valve, such as a back pressure regulator situated just upstream of the collection vessel, can be used to allow the fluid to flow into the vessel at the set pressure. An additional back pressure regulator situated at the outlet of the first collection vessel can be used to maintain the first collection vessel at a pressure lower than that of the incoming fluid.

[0058] Maintaining the first collection vessel at a pressure of between about 1000 to about 1200 psi introduces a 100 psi to 800 psi pressure drop in the collection vessel relative to the fluid entering the collection vessel. The increase in temperature and decrease in pressure creates conditions that are not conducive to keeping the kavalactones, and kavain and dihydrokavain in particular, in solution. As a result, the kavalactones precipitate out of solution and can be collected in the collection vessel.

[0059] A second and third collection vessel can be situated in series with the first collection vessel such that a generally constant pressure in the two vessels of between about 1000 psi and about 1200 psi is maintained. However the temperature of each collection vessel increases so that, for example, the second collection vessel is maintained at a temperature above that of the first collection vessel, such as between about 35° C. and about 45° C., and the third collection vessel is maintained at a temperature above that of the second collection vessel, such as between about 40° C. and about 50° C.

[0060] In a third embodiment, CO₂ is used in the supercritical phase to perform the second extraction step with the addition of ethyl alcohol as a solvent modifier. The conditions are maintained identical to that described in (1) with the addition of ethyl alcohol at about 2% to about 15% and preferably between about 2% and about 10% of ethyl alcohol content by total mass. During the process the extractable material is collected within a stainless steel collection vessel which can be the same vessel as used in the first "low pressure" preprocessing step, and the supercritical CO₂ can either be recycled for future use or vented into the atmosphere. In this embodiment the extract will contain ethyl alcohol that can be removed in subsequent processing steps. The addition of ethyl alcohol increases the polarity of the solvent, resulting in enhanced extraction of the kavalactones with higher polarity, such as methysticin and dihydromethysticin, relative to the other kavalactones in the source material.

[0061] In a fourth embodiment, the conditions are generally the same as in the third embodiment but liquid CO₂ is used instead of supercritical CO₂.

[0062] In a fifth embodiment, refrigerant chemicals, such as HFCs, HCFCs, and CFCs, are used in the liquid phase to perform the extraction step. A selection of suitable HFCs, HCFCs, and CFCs appears in Table 2. During the extraction, the refrigerant chemical(s) are maintained in the liquid state and the pressure is kept below about 400 psi. Because of the comparatively low pressure, the extraction is varied prima-

rily by temperature. Preferably, the temperature is maintained at between about 20° C. and about 70° C. and, in order to ensure suitable control and predictability, to within about $\pm 0.1^\circ$ C. of a target temperature. The refrigerant chemicals pass through the stainless steel extraction vessel containing the powdered root material and the extract-laden liquid is deposited into a stainless steel collection vessel (which can be the same one as that used in the preprocessing step). The refrigerant chemical of choice can be reclaimed via collection of the refrigerant chemical in its vapor state and subsequent pressurization into the liquid state. This is, in effect, a distillation process that leaves behind the root extract while recovering the refrigerant chemical as a liquid. The refrigerant chemical can then be re-cycled through the same mass of powdered root to repeat the process until all extractable material is removed from the powdered root or, leveraging changes in temperature and extraction solvent, until the extract has a desired kavalactone concentration.

[0063] The above-discussed processing steps will produce a kava root extract as a viscous oil, resin, or paste. According to one aspect of the invention, the preferential and/or differing rates of extraction of the various kavalactones in the feedstock for different solvents and extraction methods can be leveraged to produce a final refined kava product that has a relative kavalactone concentration meeting predefined specifications, even when those specifications may differ considerably from the concentrations present in the input kava feedstock.

[0064] Multiple extraction steps can be performed. In addition, and although not necessary, extraction can also be performed during the compression-decompression preprocessing step to selectively alter various kavalactone concentrations.

[0065] According to a particular aspect of the invention, an improved paste extract can be produced by performing an additional extraction step comprising use of supercritical CO₂ as a mobile phase combined with an adsorbent material such as that used for solid-phase extractions (SPE) and/or diatomaceous earth. The bonded materials used as silica sorbents include but are not limited to: C18 (Octadecyl), C8 (Octyl), C2 (Ethyl), CH (Cyclohexyl), PH (Phenyl), and CN (End-capped Cyanopropyl). In operation, the extract paste material obtained by the initial processing, such as via the methods discussed above, is mixed with the SPE bonded phase in a ratio of about 5:1 to about 1:1 (SPE bonded phase to root extract) by mass to produce a mixture having a doughy consistency. The mixture is then placed in an extraction vessel. Fractions can then be collected over time by pressurizing the vessel to between about 1100 psi and about 8000 psi and a temperature between about 31° C. and about 80° C. The material that elutes from this fractionation can be segregated from each other by using multiple stainless steel collection vessels. Each fraction will generally have a different percentage of kavalactones, as well as a unique distribution of kavalactones that is distinct from that seen in the original root material. Typically, the percentages of kavain and dihydrokavain are increased relative to the other kavalactones in the extract.

[0066] As will be appreciated, the time during which a solvent is allowed to flow over the kava feedstock can also effect the ratio of the final product since, for example, even a solvent that extracts a particular kavalactone slowly may

eventually extract all of that kavalactone if sufficient time has passed. The particular amount of time to allow the solvent to operate before extraction of dissolved material from the solvent begins is dependent on the kavalactone distribution of the feedstock, the different rates at which the solvent extracts each of the kavalactones, and the desired kavalactone profile of the kava extract. Given the information presented herein, one of ordinary skill in the art, using only routine experimentation can select an appropriate time to terminate the extraction process.

[0067] The composition of a kava product can be described in a variety of ways. In accordance with the present invention, several parameters are defined which can be used to describe the composition of a kava extract:

[0068] 1) Combined weight percentage of the six major alpha-pyrone: methysticin (M), dihydromethysticin (DHM), yangonin (Y), desmethoxyyangonin (DMY), kavain (K), and dihydrokavain (DHK);

[0069] 2) Combined weight percentage of M and DHM;

[0070] 3) Combined weight percentage of DHK and K;

[0071] 4) Combined weight percentage of Y and DMY;

[0072] 5) Ratio of the weight percentages of Y and DMY, i.e., Y/DMY;

[0073] 6) Ratio of the weight percentages of DHK and K, i.e., DHK/K;

[0074] 7) Combined weight percentage of flavokavain A and flavokavain B.

[0075] The first property is a rough indicator of the potency of the extract, e.g., if the extract has 0% total kavalactones it has no potency, and if it has 100% total kavalactones it is very potent. Properties 2-6 combine to define whether the kava extract can be classified in the "Daily" or "non-daily" categories, such as "Two-Day" and "No Drink". Property 7 is an indicator of the amount of flavokavains in the extract, Flavokavains are flavanoids specific to the kava plant. Flavanoids in general are antioxidants.

[0076] A most preferred kavalactone distribution for a kava extract, such as in a kava paste or a dry flowable powder for use in oral delivery has a combination of the following properties:

[0077] 1) Total weight percentage of M+DHM+Y+DMY+DHK+K ranges from a minimum of about 20% to a maximum of about 90%;

[0078] 2) Combined M+DHM weight percentage ranges from a minimum of about 15% or lower to a maximum of about 29%;

[0079] 3) Combined weight percentage of DHK and K ranges from a minimum of about 50% to a maximum of about 70%-80%;

[0080] 4) Combined weight percentage of Y and DMY ranges from a minimum of about 5% to a maximum of about 25%;

[0081] 5) Ratio of Y weight percentage to DMY weight percentage, expressed as the logarithmic function $10 \cdot \text{LOG}_{10}(Y/DMY)$ in dB units, ranges from a minimum of about -1 to a maximum of about 2;

[0082] 6) Ratio of DHK weight percentage to K weight percentage, expressed as the logarithmic function $10 \cdot \text{LOG}_{10}(DHK/K)$ in dB units, ranges from a minimum of about -4 to a maximum of about 1; and

[0083] 7) Combined weight percentage of flavokavain A and flavokavain B ranges from a minimum of about 0.3% to a maximum of about 3%.

[0084] The value of these properties for a sample of a processed kava product can be determined using conventional analytical techniques, such as HPLC-UV-S (High Performance Liquid Chromatography with Ultra-Violet detection (254 nm) and Chemical Standards), or HPLC-Electrospray-Mass Spectrometry.

[0085] Property 1 indicates that in order to have a potent product, such as a tablet, it should have a minimum of about 20% kavalactones by weight, and a maximum of about 90% kavalactones by weight. Property 7 indicates that the weight percentage of the flavokavains should be about 0.3% to about 3%, which values are typical for unprocessed kava. However, in conventional processed kava extracts, the flavokavains tend to be removed, either deliberately or due to processing effects.

[0086] Properties 2-4 combine to define extracts with a kavalactone distribution typical of a "Daily" kava cultivar found in Vanuatu. Properties Attributes 2, 5 and 6 also combine to define extracts with a kavalactone distribution typical of a "Daily" kava cultivar found in Vanuatu. The range of properties 2-7 are taken from the values measured for "Daily" kava cultivars found in Vanuatu.

[0087] According to a preferred particular embodiment, a processed kava product produced according to the invention comprises less than about 15% of the combination of methysticin and dehyromethysticin, less than about 8% of the combination of yangonin and desmethoxyyangonin, and greater than about 70% of the combination of kavain and dihydrokavain. A similar kavalactone composition comprises up to about 7% methysticin, up to about 5% dehyromethysticin, up to about 1% yangonin, up to about 4% desmethoxyyangonin, and greater than about 38% kavain.

[0088] In the extraction methods discussed above, the desired kavalactones were extracted from a source material, such as ground kava root or a pre-processed kava paste, and the remainder contained the undesirable materials. The opposite technique can also be used. In particular, a kava source material can be processed using a solvent that preferentially extracts undesired kavalactones, such as methysticin and dehyromethysticin, faster than other, more desired kavalactones. The material remaining in the extraction vessel would then be collected and could have the desired altered kavalactone profile. It has been determined that the extraction rate of methysticin and dehyromethysticin increases with increasing polarity of the extract and that certain refrigerant solvents, such as R22, can extract these undesirable kavalactones at a sufficiently greater rate than remaining kavalactones to allow this alternative refinement tech-

nique to be used. The specific extraction environments, rates of extraction, and solvent used depends on the starting profile of the source material and the degree of profile change desired. Specific solvent and environmental attributes can be determined by those of ordinary skill in the art using no more than routine experimentation typical for adjusting a process to account for, e.g., variations in the attributes of starting materials that is to be processed to produce an output material that has specified attributes.

[0089] A kava paste extract, such as discussed above, can be processed to produce consumable items, for example, by mixing it in a food product or in a capsule, or providing the paste itself for use as a dietary supplement, with sweeteners and flavors added as appropriate. However, a paste extract of this type can be difficult to process because it does not flow well and may therefore not be well suited for processing by various types of high speed machinery.

[0090] According to a further aspect of the invention, the kava paste extract discussed above (or a kava paste produced using other techniques) can be further processed to produce a dry, flowable kava powder. The powder can be used as a dietary supplement that can be added to various edible products. The powder is also suited for use in a rapid dissolve tablet.

[0091] According to a particular aspect of the invention, the kava extract powder is produced to have a kavalactone distribution that is particularly well suited for delivery in the oral cavity of human subjects, e.g., via a rapid dissolve tablet. Preferably, the desired distribution properties are achieved by adjusting the extraction process to take advantage of the differing extraction rates of the various kavalactones according to type of extraction method and the operating parameters used. Although this technique may not result in an extract that contains substantially all of the kavalactones from a sample (as is the goal of conventional extraction processes), the desired ratios can be achieved without having to perform extraction of individual kavalactones, e.g., through chromatography, and then a subsequent recombination of the individual extracts.

[0092] In one embodiment of a method for producing the kava powder, the extract paste is mixed with a suitable solvent, such as ethyl alcohol or water, along with a suitable food-grade carrier material, such as maltodextrin, dextrose, or starch and the mixture is spray air-dried using conventional techniques to produce a powder having grains of very small kava extract particles combined with the food-grade carrier material.

[0093] In a particular example, a kava extract paste, preferably about 60% to about 80% total kavalactones by weight, is mixed with about twice its weight of a food-grade carrier, such as maltodextrin having with an average particle size of between about 100 to about 150 micrometers and an ethyl alcohol solvent using a high shear mixer. Inert carriers, such as silica, preferably having an average particle size on the order of about 1 to about 50 micrometers, can be added to improve the flow of the mixture. Preferably, such additions are up to about 2% by weight of the mixture. The amount of ethyl alcohol used is preferably the minimum needed to form a solution with a viscosity appropriate for spray air-drying. Typical amounts are in the range of between about 5 to about 10 liters per kilogram of paste. The solution of kava extract, maltodextrin and ethyl alcohol is

spray dried to generate a powder with an average particle size comparable to that of the starting carrier material and a kavalactone content of about 20% to about 35% by weight.

[0094] In a second embodiment, the kava extract paste and a food-grade carrier, such as magnesium carbonate, a whey protein, or maltodextrin, are dry mixed, followed by mixing in a high shear mixer containing a suitable solvent, such as ethyl alcohol or water. The mixture is then dried via freeze drying or refractive window drying. In a particular example, kava extract paste comprising on the order of about 60% to about 80% total kavalactones by weight is combined with about one and one-half times by weight of the paste of a food-grade carrier, such as magnesium carbonate, having an average particle size of about 20 to about 100 micrometers. Inert carriers, such as silica, and preferably having an average particle size of about 1 to about 50 micrometers can be added, preferably in an amount up to 2% by weight of the mixture, to improve the flow of the mixture. The magnesium carbonate and silica are then dry mixed in a high-speed mixer, similar to a food processor-type of mixer, operating at 100's of rpm. The kava extract paste then is heated until it flows like a heavy oil. Preferably, it is heated to at least about 50° C. The heated kava paste/oil is then added to the magnesium carbonate and silica powder mixture that is being mixed in mixer.

[0095] The mixing of the kava paste/oil and the magnesium carbonate and silica is continued preferably until the particle sizes are in the range of between about 250 micrometers to about 1 mm. Between about two to ten liters of cold water (preferably at about 4° C.) per kilogram of paste is introduced into a high shear mixer. The mixture of kava extract paste, magnesium carbonate and silica is the introduced slowly or increments into the high shear mixer while mixing. An emulsifying agent, such as carboxymethylcellulose can also be added to mixture if needed. Sweetening agents can also be added at this stage if desired, such as up to about 5% by weight. Alternatively, extract of *Stevia rebaudiana*, a very sweet-tasting dietary supplement, can be added instead of, or in conjunction with, a specific sweetening agent. (For simplicity, Stevia will be referred to herein as a sweetening agent.) After mixing is completed, the mixture is dried using freeze-drying or refractive window drying. The resulting dry flowable powder of kava extract, magnesium carbonate, silica and optional emulsifying agent and optional sweetener has an average particle size comparable to that of the starting carrier material and a kavalactone content of about 20% to about 35% by weight.

[0096] According to another embodiment, kava extract paste that is on the order of about 60% to about 80% total kavalactones by weight is combined with approximately an equal weight a food-grade carrier, such as whey protein, preferably having an average particle size of between about 200 to about 1000 micrometers. Inert carriers, such as silica preferably having an average particle size of between about 1 to about 50 micrometers, or carboxymethylcellulose, preferably having an average particle size of between about 10 to about 100 micrometers, can be added to improve the flow of the mixture. Preferably, this addition is no more than about 2% by weight of the mixture.

[0097] The whey protein and inert ingredient are then dry mixed in a food processor-type of mixer that operates at 100's of rpm. The kava extract paste is heated until it flows

like a heavy oil. Preferably, it is heated to at least 50° C. The heated kava paste/oil is then added to the whey protein and inert carrier that is being mixed in the food processor-type mixer. The mixing of the kava paste/oil and the whey protein and inert carrier is continued until the particle sizes are in the range of about 250 micrometers to about 1 mm. Next, 2 to 10 liters cold water (preferably about 4C) per kilogram of paste is introduced into a high shear mixer. The mixture of kava extract paste, whey protein and inert carrier is preferably introduced slowly or in incremental amounts with the high shear mixer while mixing. Sweetening agents or other sweet-tasting additives of up to about 5% by weight, such as Stevia, can be added at this stage if desired.

[0098] After mixing is completed, the mixture is dried using freeze-drying or refractive window drying. The resulting dry flowable powder of kava extract, whey protein, inert carrier and optional sweetener has an average particle size of about 150 to about 700 micrometers and a kavalactone content of about 20% to about 35% by weight.

[0099] In a third embodiment, the kava extract paste is dissolved in a supercritical fluid, which is then adsorbed onto a suitable food-grade carrier, such as maltodextrin, dextrose, or starch. Preferably, supercritical CO₂ is used as the solvent. Specific examples include starting with a kava extract paste that is on the order of about 60% to about 80% total kavalactones by weight and adding from 1 to one one-half times the paste by weight of a food-grade carrier, such as maltodextrin, and having an average particle size of between about 100 to about 150 micrometers. This mixture is placed into a chamber containing mixing paddles and which can be pressurized and heated. The chamber is pressurized with CO₂ to a pressure in the range of between about 1100 to about 8000 psi and set at a temperature in the range of between about 20° to about 100° C. The exact temperature and pressure is selected to place the CO₂ in a supercritical fluid state.

[0100] Once the CO₂ in the chamber is in the supercritical state, the kava extract paste is dissolved. The mixing paddles agitate the carrier powder so that it has intimate contact with the supercritical CO₂ that contains the dissolved kava extract paste. The mixture of supercritical CO₂, dissolved kava extract paste and the carrier powder is then vented through an orifice in the chamber so that it undergoes explosive decompression into a collection vessel which is at a pressure and temperature that does not support the supercritical state for the CO₂. The CO₂ is thus dissipated. The resulting powder in the collection vessel is the carrier powder impregnated with the kava extract paste. The powder has an average particle size comparable to that of the starting carrier material and a kavalactone content of between about 20% to about 40% by weight. The resulting powder is dry and flowable. If needed, the flow characteristics can be improved by adding inert ingredients to the starting carrier powder, such as silica at up to about 2% by weight as discussed above.

[0101] Once a dry kava powder is obtained, such as by the methods discussed herein, it can be distributed for use, e.g., as a dietary supplement or for other uses. In a particular embodiment, the powder is mixed with other ingredients to form a tableting composition of powder which can then be formed into tablets. In a particular embodiment, the tableting powder is first wet with a solvent comprising alcohol,

alcohol and water, or other suitable solvents, in an amount sufficient to form a thick doughy consistency. Suitable alcohols include, but are not limited to, ethyl alcohol, isopropyl alcohol, denatured ethyl alcohol containing isopropyl alcohol, acetone, and denatured ethyl alcohol containing acetone. The resulting paste is then pressed into a tablet mold. An automated molding system, such as described in U.S. Pat. No. 5,407,339 can be used. The tablets are then removed from the mold and dried, preferably by air-drying for at least several hours at a temperature high enough to drive off the solvent used to wet the tableting powder mixture, typically between about 70° to about 85° F. The tablets can then be packaged for distribution.

[0102] A wide variety of tablet formulations can be made. Preferably, the tablet has a formulation that results in a rapid dissolution or disintegration in the oral cavity. The tablet is preferably of a homogeneous composition that dissolves or disintegrates rapidly in the oral cavity to release the kava extract content over a period of about 2 seconds or less to about 60 seconds or more, preferably about 3 to about 45 seconds, and most preferably between about 5 to about 15 seconds.

[0103] Various rapid-dissolve tablet formulations known in the art can be used. Representative formulations are disclosed in U.S. Pat. Nos. 5,464,632, 6,106,861, and 6,221,392, the entire contents of which are expressly incorporated by reference herein. A particularly preferred tableting composition or powder contains about 10 % to about 60% by weight of the kava extract powder and about 30% to about 60% of a water-soluble diluent. Suitable diluents include lactose, dextrose, sucrose, mannitol, and other similar compositions. Lactose is a preferred diluent but mannitol adds a pleasant, cooling sensation and additional sweetness in the mouth. More than one diluent can be used. A sweetener can also be included, preferably in an amount of between about 3% to about 40% by weight depending on the desired sweetness. Preferred sweetening substances include sugar, saccharin, sodium cyclamate, aspartame, and Stevia extract, used singly or in combination, although other sweeteners could alternatively be used. Flavorings, such as mint, cinnamon, citrus (e.g., lemon or orange), can also be included, preferably in an amount between about 0.001% to about 1% by weight.

[0104] Typically, the kava extract added to the tablet has a golden color and it is considered unnecessary to include additional colorings. However, if a coloring is desired, natural and/or synthetic colors can be added, preferably in an amount of between about 0.5% to about 2% by weight.

[0105] Typically, this tableting composition will maintain its form without the use of a binder. However, if needed, various binders are suitable and can be added in an amount of between about 5% to about 15% or as necessary. Preferred binders are acacia or gum arabic. Alternative binders include sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, hydroxyethylcellulose, methylcellulose, polyvinylpyrrolidone, VEEGUM® (available from R. T Vanderbilt Co., Inc. of Norwalk, Conn.), larch arabogalactan, gelatin, Kappa carrageenan, copolymers of maleic anhydride with ethylene or vinyl methyl ether.

[0106] A tablet according to this aspect of this invention typically does not require a lubricant to improve the flow of

the powder for tablet manufacturing. However, if it is so desired, preferred lubricants include talc, magnesium stearate, calcium stearate, stearic acid, hydrogenated vegetable oils, and carbowax in amounts of between about 2% to about 10% by weight.

[0107] Similarly, a disintegrant is not expected to be necessary to produce rapid dissolve tablets using the present tablet composition. However, a disintegrant can be included to increase the speed with which a resulting tablet dissolves in the mouth. If desired, between about 0.5% to about 1% by weight of a disintegrant can be added. Preferred disintegrants include starches, clays, celluloses, algin, gums, crosslinked polymers (including croscarmellose, crospovidone and sodium starch glycolate), VEEGUM® HV, agar, bentonite, natural sponge, cation exchange resins, alginic acid, guar gum, citrus pulp, sodium lauryl sulphate in an amount of about 0.5% to about 1% of the total mass of the tablet.

[0108] It is also generally considered unnecessary to buffer the tablet composition. However, a buffer may be beneficial in specific formulations. Preferred buffering agents include mono- and di-sodium phosphates and borates, basic magnesium carbonate and combinations of magnesium and aluminum hydroxide.

[0109] In a preferred implementation, the tableting powder is made by mixing in a dry powdered form the various components as described above, e.g., active ingredient (kava extract), diluent, sweetening additive, and flavoring, etc. An overage in the range of about 10% to about 15% of the active extract of the active ingredient can be added to compensate for losses during subsequent tablet processing. The mixture is then sifted through a sieve with a mesh size preferably in the range of about 80 mesh to about 100 mesh to ensure a generally uniform composition of particles.

[0110] The tablet can be of any desired size, shape, weight, or consistency. The total weight of the kava extract in the form of a dry flowable powder in a single oral dosage is typically in the range of about 80 mg to about 600 mg. An important consideration is that the tablet is intended to dissolve in the mouth and should therefore not be of a shape that encourages the tablet to be swallowed. The larger the tablet, the less it is likely to be accidentally swallowed, but the longer it will take to dissolve or disintegrate. In a preferred form, the tablet is a disk or wafer of about 1/8 inch to about 1/2 inch in diameter and about 0.2 inch to 0.08 inch in thickness, and has a weight of between about 160 mg to about 1,200 mg. In addition to disk, wafer or coin shapes, the tablet can be in the form of a cylinder, sphere, cube, or other shapes. For example, the tablet can be formed into the general shape of a kava plant leaf. Although the tablet is preferably homogeneous, the tablet may alternatively be comprised of regions of powdered kava extract composition separated by non-kava extract regions in periodic or non-periodic sequences, which can give the tablet a speckled appearance with different colors or shades of colors associated with the kava extract regions and the non-kava extract regions.

[0111] An exemplary 250 mg tablet contains about 125.0 mg powdered kava extract, about 12.5 mg extract of Stevia, about 35.5 mg carboxymethylcellulose, and about 77.0 mg lactose. An exemplary 350 mg tablet contains about 160.0 mg powdered kava extract, about 15.0 mg extract of Stevia,

about 15.0 mg acacia, and about 160.0 mg lactose. Other formulations are also possible.

[0112] Although the extraction techniques are discussed herein in terms of kava, it should be recognized that the disclosed techniques can be adapted for use in forming an extract, in the form of a dry flowable powder or another form and containing compositions extracted from other plant products, such as ginseng, cherry, lettuce, echinacea, mate, and areca. In addition, while the preferred rapid dissolved formulation is disclosed herein in the context of kava extract, the formulation can also be used to provide a rapid dissolve tablet containing additional or other active ingredients that can be provided in powder form, such as varieties of ginseng, cherry, lettuce, echinacea, mate, and areca.

EXAMPLES

[0113] The following examples are illustrative of the nature of the present invention and are not to be regarded as limiting.

Example 1

[0114] This example describes the preparation of a kava extract having the properties shown in FIGS. 1-4 (i.e., properties typical of a "Daily" kava cultivar from Vanuatu). This process can be varied as needed and in accordance with the profile of the source kava and the desired kava profile of the extract.

[0115] About 30 lbs. of chopped and ground lateral roots of kava are added to a 32 liter extraction vessel. Pre-cooled liquid carbon dioxide (the pressurized liquid carbon dioxide was passed through a heat exchanger that was maintained at 0° C.) was pumped through the vessel at a pressure of 1800 psi. The carbon dioxide expanded and flashed into a gas, thus cooling the lateral root to approximately 10° C. The resultant temperature equilibrium of the effluent flowing from the extraction vessel was between 5° C. and 20° C.

for "Daily", Two-Day" and "No Drink" kava cultivars from Vanuatu. The plots show the changes in the kavalactone profile introduced during processing according to the present invention. The plots show how the kavalactone profile of the processed product is shifted away from an undesirable profile to a more desired profile.

[0117] More generally, FIGS. 1-4 show how a raw kava product with a kavalactone distribution that is not typical of the preferred "Daily" kava can be processed to yield an extract with a kavalactone distribution more typical of the "Daily" kava. Note that the processing method simultaneously changes all Properties 2-6, discussed above, as well as increasing the total kavalactone weight percentage. In an optional step, the extract laden liquid carbon dioxide was collected into a cascade of three different vessels that are maintained at different temperatures and under pressures so as to further alter the kavalactone profile of the extract collected in each vessel.

Example 2

[0118] A common kava feedstock was processed and passed through sequentially connected collection vessels containing different temperature and pressure environments to produce different extraction conditions. In particular, three vessels were used containing liquid, supercritical and gaseous CO₂, respectively. The profile of the kava extracts at each stage of processing compared with the profile of the source. The first collection vessel was used for liquid CO₂ extraction. The vessel was pressurized to 1150 psi and maintained at 26.2° C. The extraction was performed at 1200 psi and 7° C. The second collection vessel was used for supercritical CO₂ extraction. The vessel was pressurized to 1150 psi and maintained at 33.5° C. The third collector was used for gaseous CO₂ collection. The collection vessel was pressurized to 850 psi and maintained between 20° C. and 30° C.

[0119] A comparison of the profile of the source feedstock and the profile of the extracts is shown in table 3, below:

TABLE 3

Kavalactone	Feedstock	CHANGE IN KAVALACTONE PROFILE (in percentages)		
		Profile in 1 st extraction vessel	Profile in 2 nd extraction vessel	Profile in 3 rd extraction vessel
Methysticin (M)	19.7	13.9	4.8	2.6
Dihydromethysticin (DHM)	10.2	10.4	13.0	8.7
Kavain (K)	22.5	41.8	39.0	28.0
Dihydrokavain (DHK)	12.7	10.9	30.0	52.4
Yangonin (Y)	22.7	5.8	1.4	0.0
Desmethoxyyangonin (DMY)	12.3	17.3	11.8	8.4

Approximately 500 lbs. of liquid carbon dioxide was passed through the lateral roots under pressure, but below the supercritical temperature. This extraction was high pressure liquid extraction. The extract laden liquid carbon dioxide was collected in a vessel that was at atmospheric pressure.

[0116] FIGS. 1-4, respectively, are plots of (DHK%+K%) vs. (M%+DHM%), (DMY%+Y%) vs. (M%+DHM%), (DMY%+Y%) vs. (DHK%+K%), and 10*LOG₁₀(DHK/K) vs. 10*LOG₁₀(Y/DMY) for the appropriate values measured

[0120] There are three dramatic changes that have occurred. First, the combined M+DHM content has dropped from 29.9% to as low as 11.3%. Second, the combined K+DHK content has increased from 35.2% to as high as 80.4%. Third, the yangonin content has dropped from 22.7% to a non-detected level, effectively 0.0%. As will be appreciated, a careful choice of the temperature and pressure for each collector will result in controlled and preferential distributions of the kavalactones.

TABLE 2

Refrigerant chemicals									
HFC							HFC mixed refrigerant		
Product name	HFC-23	HFC-32	HFC-125	HFC-134a	HFC-143a	HFC-152a	R-404A	R-407C	R-410A
Component	HFC-23	HFC-32	HFC-125	HFC-134a	HFC-143a	HFC-152a	HFC-125/143a/134a	HFC-32/125/134a	HFC-32/125
Chemical Formula	CHF ₃	CH ₂ F ₂	CHF ₂ CF ₃	CH ₂ FCF ₃	CH ₃ CF ₃	CH ₂ CHF ₂	1	2	3

HCFC				Mixed refrigerant with CFC		CFC		
Product name	HCFC-22	HCFC-123	HCFC-141b	HCFC-142b	R-502	R-11	R-12	R-113
Component	HCFC-22	HCFC-123	HCFC-141b	HCFC-142b	HCFC-22/CFC-115	CFC-11	CFC-12	CFC-113
Chemical Formula	CHC ₁ F ₂	CHC ₁₂ CF ₃	CH ₂ CC ₁₂ F	CH ₃ CC ₁ F ₂	(Azeotropic blend) 4	CC ₁₃ F	CC ₁₂ F ₂	CC ₁₂ FCC ₁ F ₂

1-CHF₂CF₃/CH₃CF₃/CH₂FCF₃(44/52/4 mass %)

2-CH₂F₂/CHF₂CF₃/CH₂FCF₃(23/25/52 mass %)

3-CH₂F₂/CHF₂CF₃(50/50 mass %)

4-CHClF₂/CC₁F₂CF₃(48.8/51.2 mass %)

1. A method of processing kava comprising the steps of:

(a) providing a kava source material having a plurality of kavalactones in a first distribution profile;

(b) extracting the kavalactones from the source material by applying a solvent to the feedstock at a predefined temperature and pressure, the solvent having different absorption rates for different kavalactones; and

(c) collecting materials dissolved in the solvent a predefined time prior to the extraction of substantially all of the kavalactones from the source material by the solvent, the predefined time selected to produce a known kavalactone ratio differential; the collected materials forming a processed kava product having a plurality of kavalactones in a second distribution ratio different from the first distribution ratio.

2. The method of claim 1, wherein the step (c) of collecting comprises collecting materials dissolved in the solvent using a plurality of series-connected collection vessels having generally constant pressure and wherein each subsequent collection vessel has a higher temperature than a preceding collection vessel, the processed kava product comprising the materials collected from at least one of the collection vessels.

3. The method of claim 1, wherein the second distribution ratio corresponds to a distribution ratio of a "one day" kavalactone cultivar.

4. The method of claim 1, wherein the combined weight percentage of methysticin, dihydromethysticin, yangonin, desmethoxyyangonin, kavain, and dihydrokavain in the processed kava product is between about 20% to about 90%.

5. The method of claim 1, wherein the combined weight percentage of methysticin and dihydromethysticin in the processed kava product is between about 5% to about 29%.

6. The method of claim 1, wherein the combined weight percentage of yangonin and desmethoxyyangonin in the processed kava product is between about 5% and about 25.

7. The method of claim 1, wherein the combined weight percentage of dihydrokavain and kavain is between about 50% to about 80%.

8. The method of claim 1, wherein LOG 10 of the ratio of the weight percentages of dihydrokavain to kavain in the processed kava product is between about -4 to about 1.

9. The method of claim 1, wherein LOG10 of a weight ratio of yangonin to desmethoxyyangonin in the processed kava product is between about -1 to about 2.

10. The method of claim 1, wherein the processed kava product comprises less than about 15% of the combination of methysticin and dehydromethysticin, less than about 8% of the combination of yangonin and desmethoxyyangonin, and greater than about 70% of the combination of kavain and dihydrokavain.

11. The method of claim 1, wherein the processed kava product comprises up to about 7% methysticin, up to about 5% dehydromethysticin, up to about 1% yangonin, up to about 4% desmethoxyyangonin, and greater than about 38% kavain.

12. The method of claim 1, wherein the processed kava product comprises a combined weight percentage of flavokavain A and flavokavain B of about 0.3% to about 3%.

13. The method of claim 1, wherein the solvent comprises liquid CO₂.

14. The method of claim 13, wherein the solvent is at a temperature lower than about 31° C. and a pressure lower than about 8000 psi.

15. The method of claim 14, wherein the solvent is at a temperature of between about 5° C. and 20° C. and the pressure is between about 1100 and about 1800 psi.

16. The method of claim 13, wherein the solvent is at a temperature of between about 5° C. and about 70° C. and a pressure below about 1100 psi.

17. The method of claim 13, wherein the solvent further comprises ethyl alcohol.

18. The method of claim 17, wherein the ethyl alcohol is added in an amount between about 2% to about 15% by mass.

19. The method of claim 18, wherein the ethyl alcohol is added in an amount between about 2% to about 10% by mass.
20. The method of claim 1, wherein the solvent comprises supercritical CO₂ at a pressure between 1100 psi and about 8000 psi and a temperature of between about 31° C. to about 80° C.
21. The method of claim 20, wherein the solvent further comprises ethyl alcohol.
22. The method of claim 21, wherein the ethyl alcohol is added in an amount between about 2% to about 15% by mass.
23. The method of claim 22, wherein the ethyl alcohol is added in an amount between about 2% to about 10% by mass.
24. The method of claim 1, wherein the solvent comprises a refrigerant chemical selected from the group consisting of hydrofluorocarbons, hydrochlorofluorocarbons, and chlorofluorocarbons.
25. The method of claim 24, wherein the solvent is maintained at a temperature between about 20° C. and about 70° C.
26. The method of claim 1, further comprising the steps of, prior to the extraction step: preprocessing the feedstock by placing it in an environment having a pressure of between about 1100 psi and about 4500 psi and rapidly reducing the pressure applied to the feedstock.
27. The method of claim 26, wherein the pressure during the preprocessing step is at least about 1800 psi.
28. The method of claim 26, wherein the temperature during the preprocessing step is between about 10° C. and about 60° C.
29. The method of claim 28, wherein the temperature during the preprocessing step is between about 15° C. and about 20° C.
30. The method of claim 26, wherein the preprocessing environment comprises CO₂.
31. The method of claim 30, further comprising the step of extracting kavalactones from the CO₂ used in the preprocessing environment.
32. The method of claim 1, wherein the source material is a kava extract, the method further comprising the step of mixing the source material with an adsorbent.
33. The method of claim 32, wherein the adsorbent material is added to the source material in an amount from about 1 times to about 5 times the amount of the source material by mass.
34. The method of claim 32, wherein the adsorbent comprises diatomaceous earth.
35. The method of claim 32, wherein the adsorbent comprises a silica sorbent.
36. The method of claim 1, wherein the source material comprises ground and dried kava plant material.
37. The method of claim 36, wherein the source material has a grain size substantially between about 50 to about 150 microns.
38. The method of claim 36, wherein the source material has a moisture content below about 7%.
39. The method of claim 1, further comprising repeating step (b) at at least one of a second pressure and a second temperature.
40. The method of claim 1, wherein the solvent is a first solvent having a first polarity;
- the method further comprising the step of repeating step (b) using a second solvent having a second polarity different from the first polarity.
41. A dry flowable kava powder prepared from a kava extract and comprising at least one kavalactone selected from the group consisting of methysticin, dihydromethysticin, yangonin, desmethoxyyangonin, kavain, and dihydrokavain.
42. The powder of claim 41, further comprising between about 0.3% and about 3% of flavokavain A and flavokavain B by combined weight of the extract.
43. The powder of claim 41, wherein the combined weight percentage of methysticin, dihydromethysticin, yangonin, desmethoxyyangonin, kavain, and dihydrokavain is between about 20% to about 90% relative to the weight of the extract.
44. The powder of claim 41, wherein the combined weight percentage of methysticin and dihydromethysticin is between about 5% to about 29% relative to the weight of the extract.
45. The powder of claim 41, wherein the combined weight percentage of yangonin and desmethoxyyangonin is between about 5% and about 25% relative to the weight of the extract.
46. The powder of claim 41, wherein LOG 10 of the ratio of the weight percentages of dihydrokavain to kavain is between about -4 to about 1.
47. The powder of claim 41, wherein LOG10 of a weight ratio of yangonin to desmethoxyyangonin in the powder is between about -1 to about 2.
48. A processed kavalactone composition comprising less than about 15% of the combination of methysticin and dehydromethysticin, less than about 8% of the combination of yangonin and desmethoxyyangonin, and greater than about 70% of the combination of kavain and dihydrokavain.
49. An ingestible kava product comprising the composition of claim 48.
50. The product of claim 49, further comprising extract of stevia.
51. A processed kavalactone composition comprising up to about 7% methysticin, up to about 5% dehydromethysticin, up to about 1% yangonin, up to about 4% desmethoxyyangonin, and greater than about 38% kavain.
52. The composition of claim 51, wherein the composition is a paste extract containing the kavalactones.
53. The composition of claim 51, wherein the composition is a liquid extract containing the kavalactones.
54. The composition of claim 51, wherein the composition is an oil containing the kavalactones.
55. The composition of claim 51, wherein the composition is a powder containing the kavalactones.
56. The composition of claim 55, further comprising a carrier.
57. The composition of claim 55, wherein the carrier is capable of absorbing lipophilic kavalactones.
58. The composition of claim 57, wherein the carrier is magnesium carbonate.
59. An ingestible kava product comprising the composition of claim 51.
60. The product of claim 59, further comprising extract of stevia.

- 61.** A powder comprising:
 about 25% to about 60% by weight of a powdered extract of kava comprising at least one kavalactone selected from the group consisting of methysticin, dihydromethysticin, yangonin, desmethoxyyangonin, kavain, and dihydrokavain; and
 about 30% to about 60% by weight of a water-soluble diluent.
- 62.** The powder of claim 61, for use as a tableting powder.
- 63.** The powder of claim 61, further comprising a lubricant.
- 64.** The powder of claim 63, wherein the lubricant is selected from a group consisting of talc, magnesium stearate, calcium stearate, stearic acid, hydrogenated vegetable oil, and carbowax.
- 65.** The powder of claim 61 for use as a capsule filling powder.
- 66.** The powder of claim 61, further comprising between about 3% and about 40% by weight of a sweetener.
- 67.** The powder of claim 66, wherein the sweetener is selected from a group consisting of sugar, saccharin, sodium cyclamate, aspartame, and extract of stevia.
- 68.** The powder of claim 61, further comprising between about 0.001% and about 1% by weight of a flavoring agent.
- 69.** The powder of claim 68, further comprising about 0.5% to about 2% by weight of a coloring agent.
- 70.** The powder of claim 61, further comprising about 5% to 15% by weight of a binder.
- 71.** The powder of claim 70, wherein the binder is selected from the group consisting of acacia, gum arabic, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, hydroxyethylcellulose, methylcellulose, polyvinylpyrrolidone, Bee-gum, alarch arabogalactan, gelatin, carrageenan, and copolymers of maleic anhydride and ethylene and/or vinyl methyl ether.
- 72.** The powder of claim 61, further comprising about 0.5% to 1% by weight of a disintegrant.
- 73.** The powder of claim 72, wherein the disintegrant is selected from a group comprising starch, clay, cellulose, algin, gum, crosslinked polymers, Veegum, agar, bentonite, natural sponge, cation exchange resins, aliginic acid, guar gum, citrus pulp, and sodium lauryl sulphate.
- 74.** The powder of claim 61, further comprising a buffer.
- 75.** The powder of claim 75, wherein the buffer is selected from a group consisting of mono- and di-sodium phosphates and borates, magnesium carbonate, and a combination of magnesium hydroxide and aluminum hydroxide.
- 76.** The powder of claim 61, further comprising a diluent.
- 77.** The powder of claim 76, wherein the diluent is selected from a group consisting of lactose, dextrose, sucrose, and mannitol.
- 78.** The powder of claim 61, wherein the combined weight of methysticin, dihydromethysticin, yangonin, desmethoxyyangonin, kavain, and dihydrokavain is between about 20% and about 90% by weight of the extract.
- 79.** The powder of claim 78, wherein the combined weight of methysticin and dihydromethysticin is between about 5% and about 29% relative to the weight of the extract.
- 80.** The powder of claim 61, wherein the combined weight of yangonin and desmethoxyyangonin is about 5% and about 25% relative to the weight of the extract.
- 81.** The powder of claim 61, wherein the combined weight of flavokavain A and flavokavain B is between about 0.3% and about 3% relative to the weight of the extract.
- 82.** The powder of claim 61, wherein LOG10 of a weight ratio of yangonin to desmethoxyyangonin in the extract is between about -1 to about 2.
- 83.** The powder of claim 61, wherein LOG10 of a weight ratio of dihydrokavain to kavain in the extract is between about -4 to about 1.
- 84.** A tablet formed from the powder of claim 61.
- 85.** A capsule filled with the powder of claim 61.
- 86.** The tablet of claim 84, wherein the powder is present in a weight of between about 80 mg to about 600 mg.
- 87.** A rapid dissolve kavalactone tablet comprising: (a) between about 25% to about 60% by weight of a powdered extract of kava; (b) between about 30% to about 60% by weight of a water-soluble diluent; (c) between about 5% to about 15% by weight of a binder; and (d) between about 3% to about 40% by weight of a sweetener, with the percentages being by weight and totaling 100%.
- 88.** The tablet of claim 87, wherein the water-soluble diluent is selected from the group consisting of lactose, dextrose, sucrose, and mannitol; wherein the binder is selected from the group consisting of acacia, gum arabic, sodium alginate, extract of Irish moss, panwar gum, ghatti gum, mucilage of isapol husks, carboxymethylcellulose, hydroxyethylcellulose, methylcellulose, polyvinylpyrrolidone, Veegum, larch arabogalactan, gelatin, carrageenan, and copolymers of maleic anhydride with ethylene and/or vinyl methyl ether; and wherein the sweetener is selected from the group consisting of sugar, saccharin, sodium cyclamate, aspartame, and extract of stevia.
- 89.** The tablet of claim 88, wherein:
 the powdered extract of kava is present in an amount of about 50% by weight;
 the diluent is lactose and is present in an amount of about 31% by weight; and
 the binder is carboxymethylcellulose and is present in an amount of about 14% by weight.
- 90.** The tablet of claim 88, wherein:
 the powdered extract of kava is present in an amount of about 46% by weight;
 the diluent is lactose and is present in an amount of about 46% by weight;
 the binder is acacia and is present in an amount of about 4%; by weight; and
 the sweetener is stevia present in an amount of about 4% by weight.
- 91.** The tablet of claim 87, wherein the tablet dissolves orally in between about 5 seconds and about 180 seconds.
- 92.** The tablet of claim 91, wherein the tablet dissolves orally in between about 15 seconds and about 60 seconds.
- 93.** A method of processing kava comprising the steps of:
 (a) providing a kava source material having a plurality of kavalactones in a first distribution profile;
 (b) extracting substantially all of the kavalactones in the feedstock using CO₂ at a first temperature and pressure to produce a kava extract having a plurality of kava-

lactones in substantially the first distribution profile, the first temperature and pressure selected to place the CO₂ in a supercritical state;

- (c) transferring the CO₂ containing dissolved kavalactones to a collection environment at a second temperature and pressure selected to cause less than all of the kavalactones dissolved in the CO₂ to precipitate out of the CO₂ within a particular time; and
- (d) collecting the precipitated kavalactones materials at the particular time, the collected materials forming a processed kava product having a plurality of kavalactones in a second distribution ratio different from the first distribution ratio, wherein at least one of the second environment and the particular time is selected produce a known difference between the first and second distribution ratios.

94. The method of claim 93, wherein the step (d) of collecting is performed using a plurality of series-connected collection vessels having generally constant pressure and wherein each subsequent collection vessel has a higher temperature than a preceding collection vessel; the processed kava product comprising the materials collected from at least one of the collection vessels.

95. The method of claim 94, wherein the generally constant pressure is between about 1000 psi and about 1200 psi and the step (d) of collection comprises collecting materials in a first collection vessel at a temperature of between about 30° C. and about 37° and a second collection vessel at a temperature of between about 35° C. and about 45° C.

96. The method of claim 95, wherein the step (d) further comprises collecting materials in a third collection vessel at a temperature of between about 40° C. and about 50° C.

97. The method of claim 93, wherein the second distribution ratio corresponds to a distribution ratio of a "one day" kavalactone cultivar.

98. The method of claim 93, wherein the second temperature is lower than about 31° C. and the second pressure is lower than about 8000 psi.

99. The method of claim 93, wherein second temperature is between about 5° C. and 20° C. and the second pressure is between about 1100 and about 1800 psi.

100. The method of claim 93, wherein the second temperature of between about 5° C. and about 70° C. and the second pressure is below about 1100 psi.

101. The method of claim 93, wherein ethyl alcohol is added to the liquid CO₂ during the extraction step (c).

102. The method of claim 101, wherein the ethyl alcohol is added in an amount between about 2% to about 15% by mass.

103. The method of claim 102, wherein the ethyl alcohol is added in an amount between about 2% to about 10% by mass.

104. A method of processing kava comprising the steps of:

- (a) providing a kava source material having a plurality of kavalactones in a first distribution profile;

- (b) removing a portion of the kavalactones from the source material by applying a solvent to the feedstock at a predefined temperature and pressure, the solvent having different absorption rates for different kavalactones; and

- (c) collecting the source material to which the solvent has been applied a predefined time prior to the removal of substantially all of the kavalactones from the source material by the solvent, the predefined time selected to produce a known kavalactone ratio differential; the collected materials forming a processed kava product having a plurality of kavalactones in a second distribution ratio different from the first distribution ratio.

105. The method of claim 104, wherein the solvent is a refrigerant.

106. The method of claim 104, wherein the plurality of kavalactones comprises a first class of kavalactones comprising methysticin and dihydromethysticin and a second class of kavalactones comprising at least one of kavain, yangonin, desmethoxyyangonin, and dihydrokavain, and wherein the solvent removes the first class of kavalactones at a greater rate than it removes the second class of kavalactones.

107. The method of claim 104, wherein the source material comprises a kava plant extract.

* * * * *